

INSTRUCTIONS FOLLOWING LASIK TREATMENT
MEDICATIONS

Day of Surgery

Pain Relief

1. Amethocaine 1% one drop as required, maximum one drop every 45 minutes. For use for today and **only if necessary** for pain relief.
2. Solpadol caps one or two 4 times a day if necessary.

Antibiotic

Maxitrol one drop: 1st dose When you wake up from snooze
2nd dose 3 hours later

Night of Surgery - Sleep

Dalmane 15 milligrams one or two on the night of treatment (Optional).

Day after Surgery

1. Continue Maxitrol 4 times a day for 6 days. Morning/Lunch/Dinner/Bed
2. Lubricating drops – Artelac drops (*preservative free*) one drop 4 hourly during the day for the next 6 weeks (*leave 10 minutes between Maxitrol and Artelac*).

INSTRUCTIONS FOLLOWING LASIK TREATMENT

1. *Remove eye shield/s on the morning following LASIK Treatment. Wear while sleeping for one week.*
2. *Do not swim for two weeks.*
3. *Do not rub eye/s for three months.*
4. *Do not wear eye make-up for one week.*
5. *Report any sudden change of vision or severe pain to the Clinic immediately.
Clinic phone lines 8.00am – 6.00pm Tel: 01-6608821/01-6688670
Emergency phone no.
(Dr. Arthur Cummings) 087-2843632
or
(Dr. Richard Corkin) 086-8234075
or
(Mr. Declan Hovenden) 086-6010510*